

**Table 10.6b**  
**Immunosuppression Use for Induction by Discharge Regimen, 2003 to 2007**  
**Recipients with Intestine Transplants**

	Discharge Regimen (w/ or w/o Steroids Use)						All
	CyA	Tac	Tac+ MMF	Tac+ Aza	Tac+ Siro	Other	
<b>Functioning Graft with Immunosuppression Info at Discharge</b>	1	509	95	4	60	27	696
<b>% of All Regimens</b>	0.1%	73.1%	13.6%	0.6%	8.6%	3.9%	100.0%
<b>Induction Drug</b>							
Atgam/NRATG/NRATS	0.0%	0.2%	4.2%	0.0%	1.7%	18.5%	1.6%
OKT3	0.0%	3.7%	30.5%	0.0%	8.3%	55.6%	9.8%
Thymoglobulin	0.0%	14.7%	51.6%	0.0%	30.0%	159.3%	26.6%
Zenapax	0.0%	11.6%	24.2%	0.0%	10.0%	81.5%	15.8%
Simulect	0.0%	2.0%	1.1%	0.0%	3.3%	14.8%	2.4%
Campath	0.0%	23.4%	3.2%	25.0%	1.7%	51.9%	19.8%
<b>No Induction Drugs Recorded</b>	100.0%	48.3%	18.9%	75.0%	55.0%	48.1%	45.1%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.