

Table 10.6d
Immunosuppression Use for Maintenance by Regimen
Prior to Discharge, 1998 to 2007
Recipients with Intestine Transplants

	Transplant Year									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Functioning Graft at Discharge	47	49	61	88	89	103	130	155	148	166
With Immunosuppression Info	47	49	61	88	89	103	127	152	148	166
Tac	0.0%	4.1%	1.6%	18.2%	31.5%	30.1%	22.8%	25.7%	16.9%	11.4%
--- + MMF/MPA	0.0%	0.0%	1.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA	0.0%	2.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.3%	7.4%	11.4%
Tac + Siro/Evero	0.0%	0.0%	0.0%	2.3%	0.0%	0.0%	2.4%	2.0%	2.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%	0.0%	0.0%
--- + Steroids	0.0%	0.0%	0.0%	1.1%	2.2%	3.9%	0.8%	2.0%	1.4%	1.8%
CyA + Steroids	6.4%	4.1%	0.0%	0.0%	0.0%	1.0%	0.0%	0.0%	0.0%	0.0%
Tac + Steroids	55.3%	67.3%	41.0%	48.9%	36.0%	50.5%	55.9%	47.4%	52.7%	56.0%
--- + MMF/MPA + Steroids	2.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	0.7%	0.0%
Tac + MMF/MPA + Steroids	27.7%	14.3%	13.1%	3.4%	7.9%	6.8%	8.7%	10.5%	10.1%	8.4%
--- + OtherAntimet + Steroids	0.0%	0.0%	1.6%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Steroids	2.1%	6.1%	0.0%	6.8%	0.0%	0.0%	0.0%	0.0%	0.7%	1.8%
--- + Siro/Evero + Steroids	0.0%	0.0%	1.6%	1.1%	0.0%	0.0%	1.6%	0.7%	0.0%	1.2%
Tac + Siro/Evero + Steroids	0.0%	0.0%	34.4%	18.2%	15.7%	3.9%	7.9%	9.9%	7.4%	6.6%
Other Regimen	6.4%	2.0%	4.9%	0.0%	6.7%	2.9%	0.0%	0.0%	0.7%	1.2%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages in each column sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.