

Table 10.6g
Immunosuppression Use for Maintenance Between Discharge
and One Year Following Transplantation, 1997 to 2006
Recipients with Intestine Transplants

	Year of Transplant									
	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Transplants	68	70	73	82	113	107	116	152	178	175
Tx with Follow-up Immunosuppression Info	31	34	37	55	69	73	88	113	130	120
Corticosteroids										
Any in Category	93.5%	97.1%	100.0%	100.0%	85.5%	90.4%	83.0%	78.8%	79.2%	86.7%
Steroids	93.5%	97.1%	100.0%	100.0%	85.5%	90.4%	83.0%	78.8%	79.2%	86.7%
Cyclosporine										
Any in Category	6.5%	2.9%	2.7%	0.0%	0.0%	1.4%	0.0%	0.9%	1.5%	1.7%
Sandimmune	0.0%	0.0%	2.7%	0.0%	0.0%	1.4%	0.0%	0.0%	0.0%	0.0%
Neoral	6.5%	2.9%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	0.8%	1.7%
Gengraf	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.8%	0.0%
Tacrolimus										
Any in Category	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	98.9%	97.3%	98.5%	98.3%
Tacrolimus	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	98.9%	97.3%	98.5%	98.3%
Antimetabolites										
Any in Category	29.0%	41.2%	37.8%	14.5%	15.9%	11.0%	10.2%	12.4%	20.0%	16.7%
Mycophenolate Mofetil	12.9%	35.3%	21.6%	14.5%	2.9%	9.6%	10.2%	8.0%	13.8%	13.3%
Azathioprine	16.1%	5.9%	16.2%	0.0%	13.0%	1.4%	0.0%	4.4%	6.2%	3.3%
mTOR Inhibitors										
Any in Category	0.0%	0.0%	21.6%	69.1%	24.6%	28.8%	15.9%	22.1%	23.8%	19.2%
Sirolimus	0.0%	0.0%	21.6%	69.1%	24.6%	28.8%	15.9%	22.1%	23.8%	19.2%
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages are calculated based on the total number of patients with any immunosuppression use between discharge and one year following transplantation. Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.