

**Table 11.6b**  
**Immunosuppression Use for Induction by Discharge Regimen, 2003 to 2007**  
**Recipients with Heart Transplants**

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
<b>Functioning Graft with Immunosuppression Info at Discharge</b>	113	257	3,362	4,503	296	116	91	45	107	827	9,717
<b>% of All Regimens</b>	1.2%	2.6%	34.6%	46.3%	3.0%	1.2%	0.9%	0.5%	1.1%	8.5%	100.0%
<b>Induction Drug</b>											
Atgam/NRATG/NRATS	8.0%	7.8%	6.1%	4.4%	7.4%	9.5%	5.5%	6.7%	4.7%	19.1%	6.5%
OKT3	2.7%	1.2%	3.4%	2.0%	25.0%	6.0%	1.1%	0.0%	2.8%	8.5%	3.7%
Thymoglobulin	6.2%	9.7%	13.4%	18.3%	18.6%	32.8%	23.1%	4.4%	41.1%	24.3%	17.2%
Zenapax	4.4%	0.8%	20.3%	10.7%	3.0%	0.0%	11.0%	77.8%	0.9%	33.5%	15.5%
Simulect	12.4%	9.7%	12.7%	14.1%	4.1%	2.6%	8.8%	6.7%	3.7%	19.5%	13.3%
Campath	0.9%	3.5%	0.1%	2.4%	0.0%	0.0%	5.5%	0.0%	0.0%	1.9%	1.5%
<b>No Induction Drugs Recorded</b>	69.0%	68.9%	46.0%	52.1%	42.9%	49.1%	46.2%	6.7%	48.6%	43.8%	49.3%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.