

Table 12.6b
Immunosuppression Use for Induction by Discharge Regimen, 2003 to 2007
Recipients with Lung Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	67	341	463	2,609	688	1,551	9	1	19	251	5,999
% of All Regimens	1.1%	5.7%	7.7%	43.5%	11.5%	25.9%	0.2%	0.0%	0.3%	4.2%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	3.0%	5.0%	0.2%	6.9%	3.9%	5.0%	0.0%	0.0%	5.3%	10.4%	5.5%
OKT3	3.0%	0.0%	1.7%	0.1%	6.3%	0.1%	0.0%	0.0%	0.0%	2.0%	1.1%
Thymoglobulin	1.5%	2.3%	2.4%	5.2%	5.8%	0.6%	11.1%	0.0%	5.3%	9.2%	3.9%
Zenapax	0.0%	2.9%	15.8%	17.7%	6.8%	16.8%	22.2%	0.0%	5.3%	40.2%	15.9%
Simulect	9.0%	6.2%	17.9%	21.8%	13.8%	32.4%	33.3%	100.0%	68.4%	53.8%	23.8%
Campath	3.0%	17.0%	0.9%	9.6%	0.0%	0.1%	0.0%	0.0%	5.3%	15.9%	6.0%
No Induction Drugs Recorded	80.6%	66.9%	61.6%	39.6%	64.2%	45.4%	33.3%	0.0%	15.8%	55.8%	48.2%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.