

Table 13.6b
Immunosuppression Use for Induction by Discharge Regimen, 2003 to 2007
Recipients with Heart-Lung Transplants

	Discharge Regimen (w/ or w/o Steroids Use)								All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	2	13	15	57	22	16	1	5	131
% of All Regimens	1.5%	9.9%	11.5%	43.5%	16.8%	12.2%	0.8%	3.8%	100.0%
Induction Drug									
Atgam/NRATG/NRATS	0.0%	7.7%	6.7%	15.8%	31.8%	0.0%	0.0%	40.0%	15.3%
OKT3	0.0%	7.7%	6.7%	1.8%	9.1%	0.0%	0.0%	0.0%	3.8%
Thymoglobulin	0.0%	0.0%	0.0%	12.3%	0.0%	12.5%	100.0%	20.0%	8.4%
Zenapax	0.0%	7.7%	26.7%	12.3%	18.2%	12.5%	0.0%	60.0%	16.0%
Simulect	0.0%	7.7%	40.0%	12.3%	13.6%	37.5%	0.0%	80.0%	20.6%
Campath	0.0%	15.4%	0.0%	21.1%	0.0%	0.0%	0.0%	60.0%	13.0%
No Induction Drugs Recorded	100.0%	69.2%	26.7%	28.1%	31.8%	37.5%	0.0%	0.0%	33.6%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.