

Table 13.6d
Immunosuppression Use for Maintenance by Regimen
Prior to Discharge, 1998 to 2007
Recipients with Heart-Lung Transplants

	Transplant Year									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Functioning Graft at Discharge	31	34	33	21	22	17	35	29	24	27
With Immunosuppression Info	29	33	33	21	22	17	35	29	24	26
CyA	3.4%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac	0.0%	0.0%	0.0%	0.0%	0.0%	5.9%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet	6.9%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Steroids	0.0%	0.0%	0.0%	4.8%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + Steroids	10.3%	12.1%	3.0%	0.0%	0.0%	0.0%	2.9%	0.0%	4.2%	0.0%
Tac + Steroids	3.4%	3.0%	3.0%	0.0%	4.5%	11.8%	11.4%	3.4%	4.2%	15.4%
CyA + MMF/MPA + Steroids	24.1%	30.3%	36.4%	33.3%	22.7%	17.6%	17.1%	10.3%	8.3%	3.8%
Tac + MMF/MPA + Steroids	10.3%	18.2%	9.1%	19.0%	18.2%	5.9%	34.3%	41.4%	62.5%	65.4%
CyA + OtherAntimet + Steroids	24.1%	18.2%	33.3%	38.1%	27.3%	23.5%	14.3%	34.5%	12.5%	0.0%
Tac + OtherAntimet + Steroids	13.8%	6.1%	6.1%	4.8%	18.2%	29.4%	14.3%	6.9%	4.2%	11.5%
Tac + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	2.9%	0.0%	0.0%	0.0%
Other Regimen	3.4%	12.1%	9.1%	0.0%	9.1%	5.9%	2.9%	3.4%	4.2%	3.8%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages in each column sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.