

Table 15.5b
Steroid Avoidance Rate at Discharge, 1998 to 2007
Recipients of Living Donor Livers

Discharge Regimen	Year of Transplant									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Total	Patients 6 8.6%	193 13 6.7%	328 30 9.1%	451 40 8.9%	308 25 8.1%	282 63 22.3%	292 64 21.9%	289 69 23.9%	260 73 28.1%	241 71 29.5%
CyA	Patients 2 25.0%	14 4 28.6%	17 1 5.9%	16 8 50.0%	6 1 16.7%	9 1 11.1%	- - -	1 0 0.0%	1 0 0.0%	1 1 100.0%
Tac	Patients 2 7.7%	96 4 4.2%	126 7 5.6%	156 11 7.1%	123 18 14.6%	101 27 26.7%	105 35 33.3%	103 36 35.0%	105 31 29.5%	70 11 15.7%
CyA+MMF	Patients 0 0.0%	14 1 7.1%	18 0 0.0%	24 0 0.0%	22 2 9.1%	18 2 11.1%	18 1 5.6%	9 2 22.2%	4 2 50.0%	7 2 28.6%
Tac+MMF	Patients 1 4.2%	50 3 6.0%	111 5 4.5%	172 2 1.2%	118 3 2.5%	136 28 20.6%	142 24 16.9%	152 23 15.1%	142 39 27.5%	142 51 35.9%
CyA+Aza	Patients 0 0.0%	1 0 0.0%	3 0 0.0%	2 0 0.0%	4 0 0.0%	3 1 33.3%	2 0 0.0%	2 0 0.0%	- - 0.0%	3 0 0.0%
Tac+Aza	Patients - - 0.0%	4 0 0.0%	2 0 0.0%	- 0 0.0%	3 0 0.0%	- - -	- - -	1 1 100.0%	- - -	2 1 50.0%
CyA+Siro	Patients - - -	- - -	7 4 57.1%	6 3 50.0%	2 0 0.0%	- - -	1 0 0.0%	- - -	- - -	- - -
Tac+Siro	Patients - - -	- - -	20 8 40.0%	36 11 30.6%	11 0 0.0%	1 0 0.0%	5 1 20.0%	1 0 0.0%	2 0 0.0%	3 0 0.0%
Siro+MMF	Patients - - -	- - -	1 0 0.0%	5 0 0.0%	2 0 0.0%	1 0 0.0%	1 0 0.0%	1 0 0.0%	- - 0.0%	2 0 0.0%
Other Regimen	Patients 1 20.0%	5 1 8.3%	12 1 21.7%	23 5 14.7%	34 5 5.9%	17 1 5.9%	13 4 30.8%	18 3 16.7%	19 7 36.8%	6 1 16.7%
										11 5 45.5%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Includes patients receiving first transplants of any organ with a functioning graft at discharge and any immunosuppressive drug use recorded at discharge.
 CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.

Table 15.5b (Continued)
Steroid Withdrawal Rate at One Year Following Transplantation, 1999 to 2006
Recipients of Living Donor Livers

Discharge Regimen	Year of Transplant							
	1999	2000	2001	2002	2003	2004	2005	2006
Total	Patients 36 Rate 24.2%	263 83 31.6%	367 164 44.7%	252 121 48.0%	195 91 46.7%	202 98 48.5%	201 102 50.7%	170 80 47.1%
CyA	Patients 1 Rate 10.0%	15 3 20.0%	7 4 57.1%	5 3 60.0%	6 2 33.3%	- - -	1 1 100.0%	1 1 100.0%
Tac	Patients 25 Rate 33.3%	101 34 33.7%	131 58 44.3%	93 43 46.2%	68 28 41.2%	65 36 55.4%	60 27 45.0%	67 28 41.8%
CyA+MMF	Patients 2 Rate 20.0%	17 4 23.5%	22 11 50.0%	16 10 62.5%	13 11 84.6%	11 7 63.6%	7 5 71.4%	2 2 100.0%
Tac+MMF	Patients 6 Rate 16.2%	94 25 26.6%	151 65 43.0%	105 55 52.4%	98 48 49.0%	108 49 45.4%	119 62 52.1%	95 49 51.6%
CyA+Aza	Patients 0 Rate 0.0%	3 1 33.3%	1 0 0.0%	4 0 0.0%	2 1 50.0%	2 0 0.0%	2 1 50.0%	- - -
Tac+Aza	Patients 1 Rate 25.0%	2 1 50.0%	- 1 50.0%	2 1 -	- - -	- - -	0 - -	- - -
CyA+Siro	Patients - - - 100.0%	3 3 2 66.7%	3 2 1 50.0%	2 1 - -	- - 100.0%	1 1 -	- - -	- - -
Tac+Siro	Patients - - - 41.7%	12 5 4 43.5%	23 10 4 44.4%	9 4 0 0.0%	1 2 2 100.0%	2 1 1 100.0%	1 1 0 100.0%	1 0 0.0%
Siro+MMF	Patients - - - 0.0%	1 0 2 50.0%	4 2 2 100.0%	2 0 0 0.0%	1 1 1 100.0%	1 1 1 100.0%	1 1 1 100.0%	- - - -
Other Regimen	Patients 1 Rate 10.0%	15 46.7%	25 48.0%	14 14.3%	6 16.7%	12 16.7%	10 40.0%	4 0.0%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Includes patients receiving first transplants of any organ with steroid use at transplant discharge, and who have a functioning graft and maintenance drug use one year after transplantation.

Transplants prior to year 1999 are not reported due to insufficient reporting of immunosuppressive drug use.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.

Table 15.5b (Continued)
Steroid Withdrawal Rate at Two Years Following Transplantation, 1998 to 2005
Recipients of Living Donor Livers

Discharge Regimen	Year of Transplant							
	1998	1999	2000	2001	2002	2003	2004	2005
Total	Patients Steroids Withdrawn Rate	47 16 34.0%	133 60 45.1%	247 132 53.4%	343 222 64.7%	236 147 62.3%	191 121 63.4%	183 121 66.1%
CyA	Patients Steroids Withdrawn Rate	5 1 20.0%	10 6 60.0%	14 9 64.3%	5 3 60.0%	5 5 100.0%	6 4 66.7%	- - 100.0%
Tac	Patients Steroids Withdrawn Rate	16 9 56.3%	63 29 46.0%	93 49 52.7%	121 75 62.0%	89 59 66.3%	64 33 51.6%	61 42 68.9%
CyA+MMF	Patients Steroids Withdrawn Rate	5 1 20.0%	7 4 57.1%	14 4 28.6%	22 17 77.3%	13 10 76.9%	13 12 92.3%	9 8 88.9%
Tac+MMF	Patients Steroids Withdrawn Rate	17 3 17.6%	37 17 45.9%	92 53 57.6%	142 98 69.0%	98 57 58.2%	99 67 67.7%	95 60 63.2%
CyA+Aza	Patients Steroids Withdrawn Rate	1 0 0.0%	3 0 0.0%	3 1 33.3%	1 0 0.0%	4 2 50.0%	2 1 50.0%	2 2 100.0%
Tac+Aza	Patients Steroids Withdrawn Rate	- - -	4 1 25.0%	2 1 50.0%	- - 100.0%	2 2 -	- - -	0 - -
CyA+Siro	Patients Steroids Withdrawn Rate	- - -	- - -	2 1 50.0%	3 3 100.0%	2 1 50.0%	- - -	1 1 100.0%
Tac+Siro	Patients Steroids Withdrawn Rate	- - -	- - -	10 7 70.0%	22 13 59.1%	9 5 55.6%	1 1 100.0%	2 2 100.0%
Siro+MMF	Patients Steroids Withdrawn Rate	- - -	- - -	1 0 0.0%	3 0 0.0%	2 2 100.0%	- - -	1 1 100.0%
Other Regimen	Patients Steroids Withdrawn Rate	3 2 66.7%	9 3 33.3%	16 7 43.8%	24 13 54.2%	12 4 33.3%	6 3 50.0%	12 5 41.7%
								10 6 60.0%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Includes patients receiving first transplants of any organ with steroid use at transplant discharge, and who have a functioning graft and maintenance drug use two years after transplantation.

Transplants prior to year 1998 are not reported due to insufficient reporting of immunosuppressive drug use.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.