

**Table 5.6f**  
**Immunosuppression Use for Maintenance by Regimen**  
**One Year Following Transplantation, 1997 to 2006**  
**Recipients with Kidney Transplants**

	Transplant Year									
	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<b>Transplants</b>	11,703	12,452	12,760	13,613	14,266	14,779	15,137	16,004	16,482	17,091
<b>Functioning Graft at 1 Year PostTx</b>	10,233	10,821	11,041	11,827	12,724	13,251	13,596	14,436	14,809	15,440
<b>With Maintenance Use Recorded at 1 Year PostTx</b>										
CyA	1,537	6,743	10,466	11,212	12,494	12,937	13,247	14,096	14,569	15,092
Tac	0.3%	0.3%	0.7%	0.6%	0.1%	0.1%	0.1%	0.1%	0.1%	0.2%
---	0.8%	0.5%	0.5%	0.5%	1.1%	1.8%	2.1%	2.2%	2.6%	2.6%
+ MMF/MPA	1.7%	0.3%	0.1%	0.0%	0.1%	0.1%	0.1%	0.2%	0.2%	0.3%
CyA + MMF/MPA	0.9%	0.9%	0.5%	1.0%	1.1%	1.5%	1.9%	2.1%	1.9%	1.8%
Tac + MMF/MPA	2.3%	1.4%	1.7%	1.8%	2.8%	5.7%	8.2%	14.1%	20.6%	24.1%
---	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
+ OtherAntimet	0.0%	0.3%	0.2%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet	0.0%	0.0%	0.2%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet	0.0%	0.1%	0.1%	0.2%	0.1%	0.1%	0.1%	0.2%	0.2%	0.4%
---	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
+ MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + MMF/MPA + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
---	0.3%	0.1%	0.1%	0.2%	0.2%	0.1%	0.1%	0.2%	0.2%	0.2%
+ Siro/Evero	0.0%	0.0%	0.2%	0.3%	0.2%	0.7%	1.0%	0.9%	0.7%	0.8%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.3%	0.2%	0.7%	1.0%	0.9%	0.7%	0.8%
Tac + Siro/Evero	0.0%	0.0%	0.1%	0.7%	1.6%	2.2%	2.4%	2.1%	1.7%	1.2%
---	0.0%	0.0%	0.0%	0.1%	0.3%	0.5%	1.1%	1.3%	1.6%	2.0%
+ MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.2%	0.0%	0.1%	0.1%	0.0%	0.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.3%	0.3%	0.4%	0.8%	0.7%	0.6%	0.5%
---	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
+ OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
CyA + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
---	2.3%	0.7%	0.5%	0.6%	0.7%	0.5%	0.4%	0.3%	0.2%	0.3%
+ Steroids	10.5%	7.8%	6.6%	4.3%	2.5%	1.6%	1.5%	1.1%	0.5%	0.4%
Tac + Steroids	6.2%	6.3%	5.9%	6.4%	5.9%	5.6%	4.9%	3.9%	3.6%	4.2%
---	0.8%	1.2%	1.5%	0.9%	0.9%	0.9%	0.8%	0.8%	1.0%	1.3%
+ MMF/MPA + Steroids	37.0%	41.5%	39.2%	31.7%	24.2%	19.9%	15.5%	12.5%	8.8%	6.5%
CyA + MMF/MPA + Steroids	17.6%	24.5%	28.3%	31.5%	37.5%	39.8%	42.6%	42.8%	43.8%	44.0%
---	0.1%	0.2%	0.2%	0.1%	0.1%	0.1%	0.1%	0.0%	0.1%	0.1%
+ OtherAntimet + Steroids	5.8%	5.6%	5.4%	3.0%	1.7%	0.9%	0.5%	0.4%	0.3%	0.2%
Tac + OtherAntimet + Steroids	2.5%	2.6%	2.2%	1.9%	2.0%	1.5%	1.3%	1.0%	0.9%	1.1%
---	0.0%	0.2%	0.3%	0.5%	0.7%	0.7%	0.6%	0.7%	0.6%	0.4%
+ Siro/Evero + Steroids	1.5%	2.5%	2.1%	4.0%	3.3%	2.7%	2.0%	1.6%	1.3%	1.0%
CyA + Siro/Evero + Steroids	0.0%	0.1%	0.9%	4.7%	7.1%	5.9%	5.1%	3.6%	2.4%	1.7%
Tac + Siro/Evero + Steroids	9.4%	2.8%	2.8%	4.4%	5.5%	6.4%	6.5%	7.2%	6.0%	4.6%
Other Regimen										

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.