

Table 5.9a
Adjusted Graft Survival by Year of Transplant at 3 Months, 1 Year,
3 Years, 5 Years and 10 Years
Deceased Donor non-ECD Kidney Transplants

	Trans- # plant Year	3 Months		1 Year		3 Years		5 Years		10 Years	
		Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.	Surv.	Std. Err.
1987	1,586	83.1%	1.0%	75.5%	1.1%	62.3%	1.2%	51.7%	1.3%	31.1%	1.2%
1988	6,858	82.7%	0.5%	75.4%	0.5%	63.1%	0.6%	53.1%	0.6%	32.1%	0.6%
1989	6,497	84.6%	0.5%	78.2%	0.5%	66.2%	0.6%	56.4%	0.6%	34.0%	0.6%
1990	6,931	86.1%	0.4%	80.0%	0.5%	68.7%	0.6%	58.1%	0.6%	35.7%	0.6%
1991	6,838	88.6%	0.4%	83.5%	0.5%	72.8%	0.5%	61.5%	0.6%	37.1%	0.6%
1992	6,653	89.3%	0.4%	84.1%	0.5%	73.1%	0.6%	61.6%	0.6%	35.7%	0.6%
1993	6,922	89.1%	0.4%	83.6%	0.4%	72.9%	0.5%	62.3%	0.6%	38.7%	0.6%
1994	6,673	90.3%	0.4%	85.2%	0.4%	75.4%	0.5%	64.0%	0.6%	39.5%	0.6%
1995	6,594	91.8%	0.3%	87.2%	0.4%	77.2%	0.5%	66.1%	0.6%	42.4%	0.6%
1996	6,523	92.9%	0.3%	88.7%	0.4%	78.7%	0.5%	67.4%	0.6%	42.9%	0.6%
1997	6,503	93.9%	0.3%	89.9%	0.4%	79.9%	0.5%	68.9%	0.6%	44.2%	0.6%
1998	6,685	94.3%	0.3%	90.6%	0.4%	81.0%	0.5%	69.9%	0.6%	+	+
1999	6,690	93.9%	0.3%	89.5%	0.4%	80.1%	0.5%	70.0%	0.6%	+	+
2000	6,788	94.1%	0.3%	89.6%	0.4%	79.5%	0.5%	68.9%	0.6%	+	+
2001	6,881	94.9%	0.3%	90.8%	0.3%	80.9%	0.5%	70.7%	0.5%	+	+
2002	7,017	95.0%	0.3%	91.1%	0.3%	81.1%	0.5%	71.3%	0.5%	+	+
2003	6,946	95.2%	0.2%	91.2%	0.3%	81.7%	0.5%	+	+	+	+
2004	7,505	95.6%	0.2%	91.9%	0.3%	82.4%	0.4%	+	+	+	+
2005	7,785	95.9%	0.2%	92.0%	0.3%	+	+	+	+	+	+
2006	8,460	96.0%	0.2%	92.4%	0.3%	+	+	+	+	+	+

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(+) = Values not determined due to insufficient follow-up.

Multi-organ transplants are excluded for kidney, pancreas, heart, lung, and liver survival rates.

Intestine survival includes intestine only and intestine with liver and/or pancreas transplants.

Living donor transplants excluded unless explicitly listed. Heterotopic heart and liver transplants excluded.

Patient survival follows patients from first transplant of this type until death. Graft survival follows individual transplants until graft failure. Counts for patient and graft survival are different because a patient may have more than one transplant for a type organ.

Survival rates are adjusted to the characteristics of transplants in year 1997. See Technical Notes for details of patient and graft survival methods.

Donors are divided into two mutually exclusive and complete categories. All donors meeting the criteria for Expanded Criteria Donors for Kidney are classified as ECD, regardless of whether the kidneys were allocated under the ECD system. Non-ECDs include all other donors, including SCDs and DCDDs.