

**Table 6.6f**  
**Immunosuppression Use for Maintenance by Regimen**  
**One Year Following Transplantation, 1997 to 2006**  
**Recipients with Pancreas Transplant Alone (PTA) Transplants**

	Transplant Year									
	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<b>Transplants</b>	79	88	139	134	167	181	160	183	199	173
<b>Functioning Graft at 1 Year PostTx</b>	46	63	109	96	122	141	115	138	148	125
<b>With Maintenance Use Recorded at 1 Year PostTx</b>	5	51	108	93	120	140	115	137	148	122
CyA	0.0%	0.0%	0.0%	1.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac	0.0%	0.0%	1.9%	2.2%	10.8%	5.7%	17.4%	15.3%	19.6%	16.4%
--- + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.0%	1.4%	9.6%	3.6%	1.4%	0.8%
CyA + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.8%	1.4%	0.9%	0.7%	0.0%	0.8%
Tac + MMF/MPA	0.0%	0.0%	3.7%	2.2%	2.5%	7.9%	7.0%	11.7%	18.2%	13.1%
Tac + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	0.7%	1.4%	0.0%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.0%	1.7%	0.7%	0.9%	2.2%	0.0%	0.0%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	1.7%	0.0%	0.7%	0.8%
Tac + Siro/Evero	0.0%	0.0%	0.9%	2.2%	0.8%	9.3%	7.0%	8.0%	8.8%	6.6%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	0.7%	1.4%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.9%	2.9%	3.4%	11.5%
Tac + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	0.0%	0.8%
--- + Steroids	0.0%	0.0%	0.0%	0.0%	0.8%	0.7%	0.0%	0.7%	0.0%	0.0%
CyA + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	0.9%	0.0%	0.0%	0.0%
Tac + Steroids	60.0%	9.8%	4.6%	9.7%	15.0%	13.6%	10.4%	12.4%	8.8%	24.6%
--- + MMF/MPA + Steroids	0.0%	0.0%	1.9%	1.1%	0.0%	1.4%	0.0%	0.0%	0.7%	0.0%
CyA + MMF/MPA + Steroids	0.0%	5.9%	5.6%	3.2%	4.2%	2.9%	2.6%	1.5%	1.4%	0.0%
Tac + MMF/MPA + Steroids	40.0%	74.5%	67.6%	52.7%	55.0%	37.9%	27.0%	22.6%	28.4%	21.3%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.7%	0.0%	0.0%
CyA + OtherAntimet + Steroids	0.0%	2.0%	0.9%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Steroids	0.0%	7.8%	0.9%	4.3%	0.8%	0.0%	0.0%	2.2%	1.4%	0.0%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	2.2%	0.0%	1.4%	0.0%	1.5%	0.0%	0.0%
CyA + Siro/Evero + Steroids	0.0%	0.0%	0.9%	0.0%	0.0%	0.7%	0.9%	0.7%	0.0%	0.8%
Tac + Siro/Evero + Steroids	0.0%	0.0%	8.3%	12.9%	6.7%	11.4%	5.2%	2.9%	4.7%	0.8%
Other Regimen	0.0%	0.0%	2.8%	6.5%	0.8%	2.1%	6.1%	8.0%	0.0%	1.6%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.