

Table 6.14
Unadjusted Patient Survival, Pancreas Transplant Alone (PTA) Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|---------------------------------|------------------|--------|-----------|------------------|--------|-----------|------------------|--------|-----------|------------------|--------|-----------|
| | (Tx 2005 - 2006) | | | (Tx 2005 - 2006) | | | (Tx 2001 - 2006) | | | (Tx 1996 - 2006) | | |
| | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 195 | 99.0% | 0.7% | 195 | 97.9% | 1.0% | 605 | 88.7% | 1.6% | 930 | 73.3% | 2.9% |
| Age at Tx | | | | | | | | | | | | |
| 6-11 Years | 0 | - | - | 0 | - | - | 0 | - | - | 1 | 100.0% | 0.0% |
| 12-17 Years | 0 | - | - | 0 | - | - | 2 | + | + | 4 | + | + |
| 18-34 Years | 54 | 100.0% | 0.0% | 54 | 98.1% | 1.8% | 161 | 85.0% | 3.7% | 269 | 71.1% | 5.2% |
| 35-49 Years | 99 | 99.0% | 1.0% | 99 | 99.0% | 1.0% | 335 | 92.1% | 1.7% | 512 | 76.0% | 3.9% |
| 50-64 Years | 42 | 97.6% | 2.4% | 42 | 95.2% | 3.3% | 104 | 81.9% | 4.9% | 141 | 64.8% | 7.9% |
| 65+ Years | 0 | - | - | 0 | - | - | 3 | 100.0% | 0.0% | 3 | + | + |
| Recipient Ethnicity/Race | | | | | | | | | | | | |
| White | 192 | 99.0% | 0.7% | 192 | 97.9% | 1.0% | 583 | 88.6% | 1.6% | 895 | 73.2% | 3.0% |
| African American | 3 | 100.0% | 0.0% | 3 | 100.0% | 0.0% | 8 | 100.0% | 0.0% | 14 | + | + |
| Hispanic/Latino | 0 | - | - | 0 | - | - | 11 | 81.8% | 11.6% | 16 | 81.3% | 9.8% |
| Asian | 0 | - | - | 0 | - | - | 0 | - | - | 1 | + | + |
| Other/Multi-race | 0 | - | - | 0 | - | - | 3 | 100.0% | 0.0% | 4 | 100.0% | 0.0% |
| Recipient Gender | | | | | | | | | | | | |
| Female | 117 | 98.3% | 1.2% | 117 | 97.4% | 1.5% | 367 | 87.1% | 2.2% | 556 | 71.1% | 4.0% |
| Male | 78 | 100.0% | 0.0% | 78 | 98.7% | 1.3% | 238 | 90.9% | 2.2% | 374 | 76.3% | 4.3% |
| Recipient Blood Type | | | | | | | | | | | | |
| O | 84 | 98.8% | 1.2% | 84 | 98.8% | 1.2% | 267 | 88.3% | 2.3% | 397 | 72.2% | 4.5% |
| A | 87 | 98.9% | 1.1% | 87 | 96.6% | 2.0% | 258 | 88.7% | 2.5% | 393 | 72.2% | 4.9% |
| B | 17 | 100.0% | 0.0% | 17 | 100.0% | 0.0% | 58 | 90.9% | 4.7% | 100 | 85.1% | 4.6% |
| AB | 7 | 100.0% | 0.0% | 7 | 100.0% | 0.0% | 22 | 88.1% | 7.9% | 40 | 61.9% | 15.5% |

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on pancreas and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Table 6.14 (Continued)
Unadjusted Patient Survival, Pancreas Transplant Alone (PTA) Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|-----------------------------|------------------|--------|-----------|------------------|--------|-----------|------------------|--------|-----------|------------------|-------|-----------|
| | (Tx 2005 - 2006) | | | (Tx 2005 - 2006) | | | (Tx 2001 - 2006) | | | (Tx 1996 - 2006) | | |
| | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 195 | 99.0% | 0.7% | 195 | 97.9% | 1.0% | 605 | 88.7% | 1.6% | 930 | 73.3% | 2.9% |
| Recipient Residency | | | | | | | | | | | | |
| U.S. Resident | 195 | 99.0% | 0.7% | 195 | 97.9% | 1.0% | 603 | 88.6% | 1.6% | 926 | 73.3% | 2.9% |
| Nonresident Alien | 0 | - | - | 0 | - | - | 2 | 100.0% | 0.0% | 2 | + | + |
| Unknown | 0 | - | - | 0 | - | - | 0 | - | - | 2 | + | + |
| Hospitalized at Tx | | | | | | | | | | | | |
| Not Hospitalized | 194 | 99.0% | 0.7% | 194 | 97.9% | 1.0% | 599 | 88.7% | 1.6% | 920 | 73.5% | 2.9% |
| Hospitalized | 1 | 100.0% | 0.0% | 1 | 100.0% | 0.0% | 6 | 83.3% | 15.2% | 9 | + | + |
| In Intensive Care | 0 | - | - | 0 | - | - | 0 | - | - | 1 | + | + |
| Cold Ischemic Time | | | | | | | | | | | | |
| 0-11 Hours | 68 | 97.1% | 2.0% | 68 | 95.6% | 2.5% | 142 | 87.4% | 3.5% | 204 | 59.7% | 9.8% |
| 12-21 Hours | 85 | 100.0% | 0.0% | 85 | 98.8% | 1.2% | 283 | 89.9% | 2.2% | 452 | 79.4% | 3.4% |
| 22-31 Hours | 16 | 100.0% | 0.0% | 16 | 100.0% | 0.0% | 60 | 88.2% | 4.6% | 103 | 62.4% | 8.6% |
| 32-41 Hours | 0 | - | - | 0 | - | - | 4 | + | + | 8 | 50.0% | 22.8% |
| 42+ Hours | 0 | - | - | 0 | - | - | 1 | 100.0% | 0.0% | 2 | 0.0% | 0.0% |
| Unknown | 26 | 100.0% | 0.0% | 26 | 100.0% | 0.0% | 115 | 88.3% | 3.6% | 161 | 81.2% | 4.2% |
| Yearly Center Volume | | | | | | | | | | | | |
| 0-4 | 7 | 85.7% | 13.2% | 7 | 85.7% | 13.2% | 19 | 100.0% | 0.0% | 55 | 84.2% | 6.8% |
| 5-7 | 8 | 100.0% | 0.0% | 8 | 100.0% | 0.0% | 29 | 96.6% | 3.4% | 72 | 82.1% | 8.4% |
| 8-11 | 25 | 100.0% | 0.0% | 25 | 100.0% | 0.0% | 67 | 93.6% | 3.1% | 68 | 76.3% | 11.1% |
| 12-19 | 22 | 100.0% | 0.0% | 22 | 100.0% | 0.0% | 97 | 93.6% | 2.8% | 104 | 77.7% | 6.4% |
| 20+ | 133 | 99.2% | 0.7% | 133 | 97.7% | 1.3% | 393 | 86.0% | 2.1% | 631 | 70.6% | 3.8% |
| Donor Age | | | | | | | | | | | | |
| <1 Year | 0 | - | - | 0 | - | - | 1 | 100.0% | 0.0% | 1 | + | + |
| 6-11 Years | 4 | 100.0% | 0.0% | 4 | 100.0% | 0.0% | 13 | 100.0% | 0.0% | 26 | 78.1% | 10.1% |
| 12-17 Years | 34 | 97.1% | 2.9% | 34 | 97.1% | 2.9% | 120 | 91.4% | 3.4% | 191 | 75.5% | 6.3% |
| 18-34 Years | 123 | 99.2% | 0.8% | 123 | 99.2% | 0.8% | 335 | 86.0% | 2.3% | 493 | 72.2% | 3.6% |
| 35-49 Years | 33 | 100.0% | 0.0% | 33 | 93.9% | 4.2% | 125 | 91.5% | 2.8% | 195 | 71.5% | 11.4% |
| 50-64 Years | 1 | 100.0% | 0.0% | 1 | 100.0% | 0.0% | 11 | 88.9% | 10.5% | 24 | + | + |

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on pancreas and kidney-pancreas transplants.

Multi-organ transplants are excluded.

Table 6.14 (Continued)
Unadjusted Patient Survival, Pancreas Transplant Alone (PTA) Transplants
Survival at 3 Months, 1 Year, 5 Years, and 10 Years

| | 3 Months | | | 1 Year | | | 5 Years | | | 10 Years | | |
|--------------------------------|------------------|--------|-----------|------------------|--------|-----------|------------------|--------|-----------|------------------|--------|-----------|
| | (Tx 2005 - 2006) | | | (Tx 2005 - 2006) | | | (Tx 2001 - 2006) | | | (Tx 1996 - 2006) | | |
| | N | % | Std. Err. |
| Total | | | | | | | | | | | | |
| All | 195 | 99.0% | 0.7% | 195 | 97.9% | 1.0% | 605 | 88.7% | 1.6% | 930 | 73.3% | 2.9% |
| Transplant Center State | | | | | | | | | | | | |
| ARIZONA | 1 | 100.0% | 0.0% | 1 | 100.0% | 0.0% | 3 | + | + | 3 | + | + |
| CALIFORNIA | 10 | 100.0% | 0.0% | 10 | 100.0% | 0.0% | 34 | 92.5% | 5.2% | 38 | 90.8% | 5.1% |
| COLORADO | 0 | - | - | 0 | - | - | 0 | - | - | 5 | 100.0% | 0.0% |
| CONNECTICUT | 1 | 0.0% | 0.0% | 1 | 0.0% | 0.0% | 4 | 75.0% | 21.7% | 11 | 62.3% | 17.9% |
| FLORIDA | 7 | 100.0% | 0.0% | 7 | 100.0% | 0.0% | 23 | 88.3% | 8.1% | 26 | 90.1% | 6.8% |
| IOWA | 0 | - | - | 0 | - | - | 5 | 100.0% | 0.0% | 9 | 88.9% | 10.5% |
| ILLINOIS | 24 | 100.0% | 0.0% | 24 | 91.7% | 5.6% | 49 | 86.6% | 7.0% | 57 | 80.8% | 9.7% |
| INDIANA | 15 | 93.3% | 6.4% | 15 | 93.3% | 6.4% | 21 | + | + | 23 | 95.7% | 4.3% |
| KANSAS | 1 | 100.0% | 0.0% | 1 | 100.0% | 0.0% | 1 | 0.0% | 0.0% | 1 | 0.0% | 0.0% |
| KENTUCKY | 0 | - | - | 0 | - | - | 0 | - | - | 1 | 0.0% | 0.0% |
| LOUISIANA | 6 | 100.0% | 0.0% | 6 | 100.0% | 0.0% | 13 | 90.0% | 9.5% | 16 | 69.2% | 20.7% |
| MASSACHUSETTS | 13 | 100.0% | 0.0% | 13 | 100.0% | 0.0% | 26 | 100.0% | 0.0% | 26 | + | + |
| MARYLAND | 11 | 100.0% | 0.0% | 11 | 100.0% | 0.0% | 50 | 85.7% | 6.3% | 124 | 68.8% | 5.6% |
| MICHIGAN | 4 | 100.0% | 0.0% | 4 | 100.0% | 0.0% | 8 | 58.3% | 25.1% | 11 | 60.6% | 18.4% |
| MINNESOTA | 38 | 100.0% | 0.0% | 38 | 100.0% | 0.0% | 197 | 85.0% | 2.8% | 309 | 60.7% | 11.1% |
| MISSOURI | 0 | - | - | 0 | - | - | 1 | + | + | 1 | + | + |
| NORTH CAROLINA | 0 | - | - | 0 | - | - | 3 | 100.0% | 0.0% | 4 | + | + |
| NORTH DAKOTA | 1 | 100.0% | 0.0% | 1 | 100.0% | 0.0% | 2 | + | + | 2 | + | + |
| NEBRASKA | 2 | 100.0% | 0.0% | 2 | 100.0% | 0.0% | 6 | 100.0% | 0.0% | 11 | 88.9% | 10.5% |
| NEW HAMPSHIRE | 1 | 100.0% | 0.0% | 1 | 100.0% | 0.0% | 1 | + | + | 1 | + | + |
| NEW JERSEY | 4 | 100.0% | 0.0% | 4 | 100.0% | 0.0% | 14 | 92.9% | 6.9% | 15 | + | + |
| NEW YORK | 2 | 100.0% | 0.0% | 2 | 100.0% | 0.0% | 7 | 85.7% | 13.2% | 19 | + | + |
| OHIO | 18 | 100.0% | 0.0% | 18 | 100.0% | 0.0% | 33 | 48.5% | 34.3% | 37 | 64.9% | 18.8% |
| PENNSYLVANIA | 12 | 100.0% | 0.0% | 12 | 100.0% | 0.0% | 38 | 92.2% | 5.6% | 54 | 92.5% | 4.2% |
| SOUTH CAROLINA | 2 | 100.0% | 0.0% | 2 | 100.0% | 0.0% | 5 | 80.0% | 17.9% | 5 | + | + |
| TENNESSEE | 4 | 100.0% | 0.0% | 4 | 100.0% | 0.0% | 10 | 100.0% | 0.0% | 35 | 59.3% | 11.4% |
| TEXAS | 2 | 100.0% | 0.0% | 2 | 100.0% | 0.0% | 7 | 100.0% | 0.0% | 15 | 75.0% | 15.8% |
| UTAH | 0 | - | - | 0 | - | - | 2 | 100.0% | 0.0% | 4 | + | + |
| VIRGINIA | 3 | 100.0% | 0.0% | 3 | 100.0% | 0.0% | 4 | + | + | 7 | 100.0% | 0.0% |
| VERMONT | 2 | 100.0% | 0.0% | 2 | 100.0% | 0.0% | 3 | + | + | 3 | + | + |
| WASHINGTON | 2 | 100.0% | 0.0% | 2 | 100.0% | 0.0% | 10 | 100.0% | 0.0% | 19 | 74.9% | 13.1% |
| WISCONSIN | 9 | 100.0% | 0.0% | 9 | 100.0% | 0.0% | 24 | 100.0% | 0.0% | 36 | 75.0% | 21.7% |

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(+) = Values not determined due to insufficient follow-up. (-) = Values not determined since there were no transplants in the category.

Cohorts are transplants performed during the years listed at the top of the table.

Patient survival follows patients from first transplant of this type until death. Counts for patient and graft survival are different because a patient may have more than one transplant for a type of organ.

Center volume = Center's yearly transplants performed during the base period, based on pancreas and kidney-pancreas transplants.

Multi-organ transplants are excluded.