

Table 7.6b
Immunosuppression Use for Induction by Discharge Regimen, 2003 to 2007
Recipients with Pancreas After Kidney (PAK) Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	5	54	63	1,030	2	6	23	23	100	158	1,464
% of All Regimens	0.3%	3.7%	4.3%	70.4%	0.1%	0.4%	1.6%	1.6%	6.8%	10.8%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.0%	0.0%	0.0%	1.7%	0.0%	0.0%	4.3%	0.0%	1.0%	1.3%	1.5%
OKT3	0.0%	0.0%	0.0%	0.4%	0.0%	0.0%	0.0%	0.0%	1.0%	1.3%	0.5%
Thymoglobulin	100.0%	20.4%	65.1%	56.9%	50.0%	50.0%	60.9%	91.3%	45.0%	108.2%	61.3%
Zenapax	0.0%	3.7%	6.3%	6.1%	0.0%	0.0%	0.0%	0.0%	7.0%	11.4%	6.4%
Simulect	0.0%	3.7%	12.7%	6.0%	0.0%	0.0%	4.3%	8.7%	3.0%	5.7%	5.9%
Campath	20.0%	46.3%	14.3%	9.7%	0.0%	33.3%	21.7%	0.0%	24.0%	55.1%	17.3%
No Induction Drugs Recorded	0.0%	29.6%	12.7%	22.7%	50.0%	16.7%	26.1%	4.3%	24.0%	8.9%	20.8%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.