

Table 8.6b
Immunosuppression Use for Induction by Discharge Regimen, 2003 to 2007
Recipients with Kidney-Pancreas Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	1	159	109	3,176	1	18	82	147	278	319	4,290
% of All Regimens	0.0%	3.7%	2.5%	74.0%	0.0%	0.4%	1.9%	3.4%	6.5%	7.4%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.0%	1.9%	0.9%	2.1%	0.0%	0.0%	1.2%	0.0%	0.4%	2.2%	1.9%
OKT3	0.0%	1.3%	0.0%	0.5%	0.0%	0.0%	1.2%	0.0%	1.4%	2.5%	0.7%
Thymoglobulin	100.0%	28.9%	57.8%	50.8%	0.0%	55.6%	19.5%	83.0%	47.1%	79.3%	52.5%
Zenapax	0.0%	5.7%	4.6%	6.9%	0.0%	5.6%	1.2%	0.0%	2.5%	4.7%	6.0%
Simulect	0.0%	2.5%	25.7%	10.7%	0.0%	33.3%	2.4%	0.7%	1.4%	8.2%	9.6%
Campath	0.0%	31.4%	0.9%	9.8%	0.0%	0.0%	69.5%	0.7%	25.9%	25.4%	13.3%
No Induction Drugs Recorded	0.0%	31.4%	13.8%	21.8%	100.0%	5.6%	8.5%	16.3%	24.1%	9.7%	20.7%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.