

Table 9.6b
Immunosuppression Use for Induction by Discharge Regimen, 2003 to 2007
Recipients with Liver Transplants

	Discharge Regimen (w/ or w/o Steroids Use)										All
	CyA	Tac	CyA+ MMF	Tac+ MMF	CyA+ Aza	Tac+ Aza	Siro+ MMF	CyA+ Siro	Tac+ Siro	Other	
Functioning Graft with Immunosuppression Info at Discharge	330	8,930	1,154	15,600	110	219	228	163	318	1,412	28,464
% of All Regimens	1.2%	31.4%	4.1%	54.8%	0.4%	0.8%	0.8%	0.6%	1.1%	5.0%	100.0%
Induction Drug											
Atgam/NRATG/NRATS	0.3%	0.6%	0.5%	0.4%	0.0%	0.0%	0.0%	0.0%	0.6%	2.0%	0.6%
OKT3	0.3%	0.7%	0.7%	2.0%	0.0%	0.9%	0.9%	0.0%	2.2%	2.5%	1.5%
Thymoglobulin	4.5%	7.7%	7.1%	7.6%	16.4%	15.1%	11.8%	2.5%	18.2%	21.2%	8.5%
Zenapax	7.6%	3.5%	9.7%	6.0%	0.0%	3.7%	8.8%	47.2%	3.1%	15.2%	6.0%
Simulect	7.9%	4.1%	19.2%	6.2%	1.8%	44.7%	11.8%	4.9%	5.7%	17.6%	7.0%
Campath	0.3%	3.0%	0.3%	0.6%	0.0%	0.9%	0.0%	0.0%	0.9%	2.3%	1.4%
No Induction Drugs Recorded	79.7%	80.9%	63.2%	79.3%	81.8%	34.7%	67.5%	46.0%	72.0%	76.6%	78.3%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Individual induction drug percentages will not necessarily add up to 100 percent, as patients may be prescribed more than one induction drug.

CyA: Cyclosporine; Tac: Tacrolimus; MMF: Include MMF(Mycophenolate Mofetil) and MPA(Mycophenolate Sodium); Aza: Azathioprine; Siro: Sirolimus.